## ERRATUM

In the October 2020 issue of *Continuum* (Peripheral Nerve and Motor Neuron Disorders, Vol. 26, No. 5), the following error occurred:

In "Chronic Inflammatory Demyelinating Polyradiculoneuropathy and Its Variants" by Kelly Gwathmey, MD (Continuum: Lifelong Learning in Neurology 2020:26:1221), the dosage for prednisone is stated incorrectly as "Several corticosteroid regimens are used for induction therapy, including oral prednisone (1 g/kg/d to 1.5 g/kg/d or alternate-day equivalent dose), dexamethasone 40 mg/d for 4 days every 4 weeks, or IV methylprednisolone 0.5 g one day each week or for 4 consecutive days monthly." The corrected dosage is shown below:

Several corticosteroid regimens are used for induction therapy, including oral prednisone (1 mg/kg/d to 1.5 mg/kg/d or alternate-day equivalent dose), dexamethasone 40 mg/d for 4 days every 4 weeks, or IV methylprednisolone 0.5 g one day each week or for 4 consecutive days monthly.

Gwathmey K. Chronic inflammatory demyelinating polyradiculoneuropathy and its variants. Continuum (Minneap Minn) 2020;26(5, Peripheral Nerve and Motor Neuron Disorders):1205-1223. doi:10.1212/CON.000000000000907